
COMPARISON OF 2-ALKYL IMMIDASOLIUM AND HALO IMMIDASOLIUM

P.Vishwamithran¹, Gurumoorthy², R.Shanmugapriya³, V.Bhakyajothi⁴

Assistant professor

¹ Department of chemistry, Dhanalakshmi Srinivasan College of arts and science for women, Perambalur, Tamil Nadu

Abstract

The molecular structures of imidazole, benzimidazole and its 2-alkyl derivatives as well as their vibrational frequencies have been determined by DFT method using B3LYP theory level at 6-311G (d,p) basis set. Some physical properties such as total energy, HOMO, LUMO energies and dipole moments of studied molecules were calculated also determined. 2-ethyl derivative possesses the lowest energy -458 hartree compare with imidazole -226.27 hartree whereas the smallest energy gap between HOMO and LUMO is in 2-methyl derivative it is the most chemically reactive. Imidazole and benzimidazole are planar molecules and have dipole moments 3.7285 and 3.5528 D respectively. C4 atom carries the highest negative charge compare with the C2 atom in imidazole. In all studied molecules N1 carries higher negative charge than N3, but N1 and N3 atoms show the highest negative charge in 2-ethyl derivative. The hydrogen atom of N-H group is coplanar in all studied molecules.

Vibration frequencies and simulated spectra have been discussed and compared with the measured spectra.

Keywords: DFT calculations, molecular structures, vibrational frequencies, imidazole, benzimidazole and its 2-alkyl derivatives.

Introduction

Imidazole and benzimidazole are aromatic heterocyclic compounds occur widely as essential constituent in the most versatile binding sites in protein. Their derivatives are


present in antibacterial, antifungal, antiprotozoal and antihelminthic medications (Katritzky et al. 2000)(DeLuca 2006) , metal ion complexes containing imidazole rings are used as metal corrosion inhibitors(otacicandStupinisck-Lisca 2003) and potential determined by MP2/6-311+G(d,p) anticancer agent(Zhoo and Lin 2005). calculations(Xu Wang et al. 2006) .

Imidazole and benzimidazole are expected to the aim of the present study is to investigate function as aromatic π ligand or simple σ the molecular structures and vibrational ligand in binding with aromatic or molecular species (Sundberg and Martin 1974). For metal complexes, binding energies of a series of frequencies of imidazole, benzimidazole and 2-alkylbenzimidazole derivatives.

Experimental and computational method of metal ions of univalent have been The studied molecules are represented with measured by collision-induced dissociation (atoms numbering by the following Figure.

Rannulu and Rodgers 2005). The metal ions in these complexes have been predicated to be σ -bound to N3 group of imidazole(Alcami and Janez 1992). Metal complexes of heterocyclic compounds have been studied by They were prepared by general method from condensation of o-phenelene diamine with corresponding aliphatic carboxylic acids. The compounds were characterized by melting points and IR spectra(Seka and Müller 1931)(zero electron kinetic energy spectroscopy(Pool et al. 1977). Imidazole was obtained

Wang et al. 2006). Structure of imidazole and from Fluka Company and used without Al-imidazole, Cu-imidazole have been further purification.



These compounds have been to liberate from showed that they are also acidic ($p^{Ka} = 5.4$) acidic solution by addition of ammonia which indicates that these heterocyclic aromatic compounds are basic. Experimental tests and stable to the hydrolytic action of alkalis, as well as that of acids (Fisser and Willimson 1975).

IR measured by SHIMADZU FTIR as KBr disc in the range 4000-500 cm^{-1} . The theoretical calculations were carried out by personal computer type Pentium-4 with CPU

GHz and ram 512 Mbit. Software package G03W program has been used which involves ab initio, DFT and semi empirical calculations. DFT molecular orbital calculations have been performed using B3LYP theory level at 6-311G (d,p) basis set. The geometry optimization was performed before the calculations of vibrational frequencies, then simulated IR spectra were obtained.

Results and discussion

Some of the calculated physical properties of studied molecules such as total energy, HOMO and LUMO orbital energies and dipole moments are shown in Table (1), whereas Mulliken charges on the atoms are listed in Table (2).

Table (1): Some physical properties of studies molecules determined by DFT method using B3LYP/6-311G (d,p) theory level-basis set.

Molecules	Total energy	HOMO energy	LUMO Energy	ΔE	Dipole moment
Imidazole	-226.226	-0.234	0.021	0.255	3.7285
Benzimidazole	-379.960	-0.232	0.012	0.244	3.4799
2-methylbenzimidazol	-419.294	-0.018	0.015	0.033	3.4177
2-ethylbenzimidazole	-458.618	-0.228	0.014	0.242	3.5528

Dipole moment in Debye unit (D), Energy in hartree unit.

It's clear from Table (1) that 2-ethylbenzimidazole has the lowest total energy which suggests it is relatively more thermally stable than the other molecules. However the energy difference between HOMO and LUMO energies shows the lowest value in 2-methylbenzimidazole which reflects its relatively high chemical reactivity compared with other studied molecules. On the other hand, the results of dipole moment showed only two components which indicate the planarity of the molecules.

However, imidazole showed a higher dipole moment value than benzimidazole which is in

agreement with the high melting point that can be explained in terms of presence of strong intermolecular interactions and hydrogen bonding. On the other hand, substitution at 2-position affects the dipole moment.

The theoretically calculated Mulliken charges on the atoms are shown in Table (2). It is clear from Table (2) that the carbon atom C4 carries the highest negative charge which suggests the electrophonic attack is more likely to occur at this such as chlorination, nitration and sulphonation in imidazole. However, nitrogen atom N1 carries the largest negative charge in all studies molecule (see Table 2). It has -0.461 in 2-ethylbenzimidazole compare with -0.328 in imidazole. In addition N1 always shows a higher negative charge than that on N3.

Table (2): Mulliken charges on some atoms determined by DFT using B3LYP theory level at 6-311G(d, p) basis set

Molecules Atom No.	Imidazole	Benz- imidazole	2-Methylbenz- imidazole	2-Ethylbenzi- midazole
N1	-0.328	-0.423	-0.454	-0.461
C2	0.152	0.1177	0.245	0.310
N3	-0.298	-0.316	-0.332	-0.345
C4	-0.073	-0.015	-0.012	-0.011
C5	0.000	0.178	0.186	-0.189
H6	0.227	0.230	0.227	0.226
H7	0.111	0.111	-----	-----
C8	-----	-0.053	-0.064	-0.056
C9	-----	-0.045	-0.053	-0.054
C10	-----	-0.011	-0.112	-0.112

Some of the geometrical parameters, bond lengths and bond angles, calculated by DFT method at B3LYP/6-311G (d,p) theory level / bases set are listed in Tables (3) and (4).

From Table (3), the two bonds, C2-N1 and C5-N1, are quite different in imidazole molecule whereas they are similar in benzimidazole, 2-methylbenzimidazole molecules. The calculated bond lengths in imidazole are in good agreement with previously reported values in the literature (Hus and Craven 1974) (Wang et al. 2006) (Kok et al. 1975). On the other hand, these bonds become longer in benzimidazole and the substituted benzimidazole. These facts

can be interpreted in term of inductive effect of the methyl and ethyl groups or the fused aromatic ring. In addition the N-H bond is unaffected by the substituent at 2-position but C3-N3 bond becomes slightly shorter while C4-N3 bond get longer than in imidazole.

From Table (4), the aromatic hydrogen atoms and N-H atom are coplanar in all studied molecules. The calculated bond angles are in agreement with previously reported values determined by X-ray diffraction measurements (Makino et al. 1999) and that calculated by MP2/6-311G + (d, p) in imidazole (Lide and Frederikse 1997).

The vibrational frequencies and their relative intensities of the studied molecules have been determined by DFT molecular orbital calculations. These molecules show 3N-6 normal modes of vibrations, the numbers of normal moods are 21, 39, 48 and 57 for imidazole, benzimidazole, 2-methylbenzimidazole and 2-ethylbenzeimidazole respectively.

Table (3): Some bond lengths (Å) of studied molecules determined by DFT method at B3LYP/ 6-311G (d,p) bases set.

molecule Bond	Imidazole	Benzimidazole	2-Methylbenzimidazole	2-ethylbenzimidazole
C2-N1	1.366	1.377	1.383	1.383
C5-N1	1.379	1.383	1.384	1.384
N1-H6	1.007	1.007	1.007	1.007
C2-N3	1.312	1.304	1.308	1.307
C2-H7	-----	1.081	-----	-----
C2-C7	-----	-----	1.492	1.500
C4-N3	1.377	1.389	1.388	1.389
C4-H8	1.079	-----	-----	-----
C4-C8	-----	1.398	1.397	1.397
C5-H9	1.077	-----	-----	-----
C5-C9	-----	1.395	1.393	1.397
C9-C10	-----	1.390	1.391	1.391
C9-C11	-----	1.084	1.084	1.084
C10-C11	-----	1.408	1.406	1.406

The vibrational frequencies between 3656 and 3724 cm^{-1} is assigned to N-H stretching (Table 5) as also shown in simulated spectra Figs (1) and (2). This band appears very broad in the solid state (KBr disk) due to strong hydrogen bonding. The vibrational frequencies at 3270 cm^{-1} in imidazole and at 3228 cm^{-1} in benzimidazole are arisen from stretching vibration of the hydrogen at 2- position (Table 5) and Fig 1. The other frequencies 3241 and 3237 cm^{-1} are attributed to C-H stretching of the heteroaromatic ring. However the four frequencies at 3196, 3186, 3175 and 3165 cm^{-1} are assigned to the stretching vibration of the aromatic hydrogen's, symmetric and antisymmetric. In addition the four frequencies at 3204, 3188, 3173 and 3163 cm^{-1} are attributed to the C-H stretching of the aromatic hydrogen in 2- methylbenzimidazole and 2- ethylbenzimidazole respectively.

Table (4): Some of the calculated bond angles in studied molecules (Angles $^{\circ}$) determined by DFT method using B3LYP /6-311G (d,p) theory bases set.

Molecule Angle	Imidazole	Benzimidazole	2-Methylbenzimidazole	2-ethylbenzimidazole
N1-C5-H9	123.3	-----	-----	-----
C4-C5-C9	-----	133.1	133.2	133.2
C2-N3-C4	105.4(105.2) ^a	104.8	105.5	105.6
N3-C2-H7	125.9	125.2	-----	-----
C2-N1-C5	107.2	106.8	107.4	107.4
N3-C4-C5	110.7(110.1) ^a	110.4	110.3	110.2
N3-C4-H8	121.4	-----	-----	-----
N3-C4-C8	-----	129.9	130	130
C4-C4-H8	127.9	-----	-----	-----
C5-C4-C8	-----	119.7	119.7	119.7
C4-C5-H9	132.6	-----	-----	-----
C4-C8-C11	-----	118.1	118.1	118.1
C4-C8-H15	-----	120.1	120.1	120.1
C5-C9-C10	-----	116.7	116.8	116.8
C5-C9-H12	-----	122.1	122.1	122.0
C2-N1-H6	126.6	126.4	126.1	126.1
N1-C2-N3	111.7(111.4) ^a	113.6	112.5	112.4
N1-C2-H7	122.4	121.3	-----	-----
N1-C2-C7	-----	122.0	121.5	-----
C5-N1-H6	126.2	126.8	126.5	126.5
C4-C5-N1	105.1(106.3) ^a	104.4	104.3	104.3
C5-N1-C2	107.2(107.6) ^a	107.2	107.2	107.3

a) Ref, K. Makino, H. S. Kim and Y. K. urasawa, J. Hetrocycl. Chem, 35, 489, (1999).

b) Ref, D. R. Lide, H. P. R. Frederiks, Handbook of Chemistry and Physics, 78 th ed.(1997).

From comparison of the data of benzimidazole and 2-methyl derivative it is clear that the frequencies at 3308, 3302, 3228 cm^{-1} , and 3132, 3113, 3096, 3035, 3033 cm^{-1} in 2-methylbenzimidazole and 2-ethylbenzimidazole respectively are attributed to the C-H aliphatic asymmetric and symmetric stretching vibrational frequencies (Figs (1) and (2)). The other frequencies between 1558-1434 cm^{-1} and 1664-1496 cm^{-1} and 1775-1445 cm^{-1} and 1661-1496 cm^{-1} in imidazole, benzimidazole, 2-methylbenzimidazole and 2-ethylbenzimidazole respectively are assigned to C=N and C=C in the aromatic rings, Table (5). These are in agreement with the experimentally measured spectra.

On the other hand, the frequencies between 1455-1384 cm^{-1} and 1477-1370 cm^{-1} in 2-methyl and 2-ethylbenzimidazole respectively are attributed to C-H bending deformation of the aliphatic groups.

From the theoretical data and the simulated spectra Figs 1 and 2, the strong frequencies at 816 and 730 cm^{-1} in imidazole, and 754 cm^{-1} benzimidazole, and 726,673 cm^{-1} in 2-ethyl, and 749,725 cm^{-1} in 2-ethylbenzimidazole could be raised from C-H out of plane bending of the aromatic hydrogen atoms whereas the frequencies between 1100-1019 cm^{-1} region may be attributed to C-H bending in plane for all the studied molecules.

From comparison of the data in Table 5, it is clear that lower frequencies at 132 and 124 cm^{-1} in 2-methylbenzimidazole are raised from torsional motions of the methyl group whereas the frequencies at 211, 181, 98 and 39 cm^{-1} are attributed to torsional motion of the ethyl group, as shown in the simulated spectra Fig (2). The corresponding peaks have not been observed in the measured spectra in the solid state since the instrument range is limited down to 500 cm^{-1} .

Table (5): Some of the theoretically calculated vibrational frequencies of the studied molecules by DFT method using B3LYP theory level at 6-311G (d,p) basis set.

Imidazole freq./intensity%		Benzimidazole freq./intensity%		2-Methylbenzimidazole freq./intensity%		2- Ethylbenzimidazole freq./intensity%	
3656	50.01	3660	63.22	3724	14.82	3645	45.92
3270	1.95	3228	2.32	3328	8.96	3204	12.18
3241	0.46	3196	11.18	3326	6.39	3118	23.59
3237	0.95	3186	22.23	3322	14.11	3173	13.68
1558	21.69	3175	12.79	3316	10.11	3164	1.97
1500	17.14	3165	0.07	3308	3.18	3132	16.69
1434	14.02	1664	6.54	3302	0.22	3113	28.47

1363	6.33	1623	3.63	3228	15.01	3096	28.93
1281	0.37	1534	25.65	1755	8.51	3035	29.74
1162	4.73	1521	4.65	1596	2.59	3033	27.52
1145	2.68	1477	25.15	1524	59.35	1661	5.66
Imidazole freq./intensity%		Benzimidazole freq./intensity%		2-Methylbenzimidazole freq./intensity%		2- Ethylbenzimidazole freq./intensity%	
1092	20.36	1421	30.48	1455	9.89	1694	0.56
1074	38.84	1378	29.46	1443	17.14	1566	65.41
964	2.05	1334	9.21	1384	11.14	1561	0.22
909	8.88	1286	30.24	1332	6.92	1501	10.32
870	4.26	1274	1.84	1326	2.97	1496	4.15
816	33.82	1203	1.04	1310	1.59	1477	3.16
730	36.18	1170	1.72	1260	44.63	1773	31.88
683	5.07	1130	2.39	1184	1.61	1443	44.95
646	11.39	1095	18.01	1138	8.95	1407	3.67
528	95.13	1028	5.36	1117	4.37	1400	29.90
		985	0.12	1086	4.29	1370	24.58
		947	1.64	1032	12.41	1332	13.04
		945	2.62	990	2.84	1293	12.04





Figure 1.


Simulated spectra calculated by DFT using B3LYP/6-311G(d,p) basis set

a-Imidazole b- benzimidazole

(1)



a



b

Figure : (2)

Simulated spectra calculated by DFT using B3YLP/6-311G(d,p) basis set

a- 2-methylbenzimidazole b- 2-ethylbenzimidazole

References

- M. Alcami, O. Mo, M. J. Janez, *J. Pys. Chem*, 96, 3012, (1992).
- L. DeLuca, *Curr. Med. Chem.* 1,13, (2006).
- L. F. Fisser and K. L. Willimson, "Organic experiments" 3rd. ed. (1975).
- I. N. Hus and B. M. Craven, *Acta Crystallogr.* 30, 988 (1974).
- A. R. Katritzky, A. F. Pozharskii, *Handbook of Hetrocyclic Chemistry*, 2nd ed, Pergamon: Amsterdam, (2000).
- A. J. Kok, C. Romas and J. Hoogendrop, *Acta Cryst*, 31, 2818 (1975).
- D. R. Lide, H. P. R. Frederikse. *CRC Handbook of Chemistry and Physics*, 78th ed. CRC: New York (1997).
- K. Makino, H. S. Kim and Y. Kuarasawa, *J. Hetrocycl. Chem*, 35,489(1999).
- H. Otacic, E. Stupinisck-Lisca, *Electrochim Acta*,48,985,(2003).
- W. O. Pool, H. J. Harwood, and A.W. Ralston, *J. Am.Soc*, 59, 178, (1977).
- S. N. Rannulu, M.T. Rodgers, *Phys. Chem. Chem. Phys.*7,1014, (2005).
- R. Seka and R.H.Müller, *Monatsh*,57,97,(1931).
- R. J. Sundberg, R. B. Martin, *Chem.Rev.*74, 471,(1974).
- X. Wang, J. S. Lee, D. S. Yang, *J. Chem. Phys*,125,14309, (2006).
- Xu Wang, J. S. Lee, Dong-Sheng Yang, *J. Phys.Chem. A*,110,12772(2006).
- X. Wang, J. S. Lee and D.S. Yang, *J. Phes.Chem. A*, 110, 5 (2006).
- G. Zhoo, H. Lin, *Curr. Med. Chem: Anti- cancer Agent*, 5, 137,(2005).